

Activin Receptor Type IIB Rabbit pAb

db20325

Package : 20µL 50µL 100µL

Product Name : Activin Receptor Type IIB Rabbit pAb**Cat.No.:** db20325**Synonyms** : HTX4; ACTRIIB; ActR-IIB**Application** : WB**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases. This gene encodes activin A type IIB receptor, which displays a 3- to 4-fold higher affinity for the ligand than activin A type II receptor. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human Activin Receptor Type IIB/ACVR2B

Gene ID

93

Swiss Prot

Q13705

Synonyms

HTX4; ACTRIIB; ActR-IIB

Reactivity

Human, Mouse, Rat

Application

WB

Recommended dilution

WB: 1:1000

Calculated MW

58 kDa

Observed MW

58 kDa

Host species

Rabbit

Clonality

Polyclonal

Isotype

IgG

Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.