

**Bcl10 Rabbit pAb**

db20523

Package : 20µL 50µL 100µL

**Product Name** : Bcl10 Rabbit pAb**Cat.No.:** db20523**Synonyms** : CLAP; mE10; CIPER; IMD37; c-E10; CARMEN**Application** : WB, IHC, ICC/IF, IP**Reactivity** : Human**Host species** : Rabbit**Background**

This gene was identified by its translocation in a case of mucosa-associated lymphoid tissue (MALT) lymphoma. The protein encoded by this gene contains a caspase recruitment domain (CARD), and has been shown to induce apoptosis and to activate NF-kappaB. This protein is reported to interact with other CARD domain containing proteins including CARD9, 10, 11 and 14, which are thought to function as upstream regulators in NF-kappaB signaling. This protein is found to form a complex with MALT1, a protein encoded by another gene known to be translocated in MALT lymphoma. MALT1 and this protein are thought to synergize in the activation of NF-kappaB, and the deregulation of either of them may contribute to the same pathogenetic process that leads to the malignancy. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016]

**Immunogen**

Recombinant protein of human Bcl10

**Gene ID**

8915

**Swiss Prot**

O95999

**Synonyms**

CLAP; mE10; CIPER; IMD37; c-E10; CARMEN

**Reactivity**

Human

**Application**

WB, IHC, ICC/IF, IP

**Recommended dilution**

WB: 1:1000

IHC: 1:50

ICC/IF: 1:50

IP: 1:20

**Calculated MW**

26 kDa

**Observed MW**

26 kDa

**Host species**

Rabbit

Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.