

Thyroid Transcription Factor 1 Rabbit pAb

db21727

Package : 20µL 50µL 100µL

Product Name : Thyroid Transcription Factor 1 Rabbit pAb**Cat.No.:** db21727**Synonyms** : BCH; BHC; NK-2; TEBP; TTF1; NKX2A; NMTC1; T/EBP; TITF1; TTF-1; NKX2.1**Application** : WB, IHC, ICC/IF, FC, IP**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

This gene encodes a protein initially identified as a thyroid-specific transcription factor. The encoded protein binds to the thyroglobulin promoter and regulates the expression of thyroid-specific genes but has also been shown to regulate the expression of genes involved in morphogenesis. Mutations and deletions in this gene are associated with benign hereditary chorea, choreoathetosis, congenital hypothyroidism, and neonatal respiratory distress, and may be associated with thyroid cancer. Multiple transcript variants encoding different isoforms have been found for this gene. This gene shares the symbol/alias 'TTF1' with another gene, transcription termination factor 1, which plays a role in ribosomal gene transcription. [provided by RefSeq, Feb 2014]

Immunogen

A synthetic peptide of human TTF1

Gene ID

7080

Swiss Prot

P43699

Synonyms

BCH; BHC; NK-2; TEBP; TTF1; NKX2A; NMTC1; T/EBP; TITF1; TTF-1; NKX2.1

Reactivity

Human, Mouse, Rat

Application

WB, IHC, ICC/IF, FC, IP

Recommended dilutionWB: 1:1000
IHC: 1:50
ICC/IF: 1:50
FC: 1:20-1:100
IP: 1:20**Calculated MW**

39 kDa

Observed MW

39 kDa

Host species

Rabbit

Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.