

PYK2 Rabbit pAb

db22335

Package : 20µL 50µL 100µL

Product Name : PYK2 Rabbit pAb**Cat.No.:** db22335**Synonyms** : PKB; PTK; CAKB; FAK2; PYK2; CADTK; FADK2; RAFTK**Application** : WB, IHC, IP**Reactivity** : Human, Rat**Host species** : Rabbit**Background**

This gene encodes a cytoplasmic protein tyrosine kinase which is involved in calcium-induced regulation of ion channels and activation of the map kinase signaling pathway. The encoded protein may represent an important signaling intermediate between neuropeptide-activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. The encoded protein undergoes rapid tyrosine phosphorylation and activation in response to increases in the intracellular calcium concentration, nicotinic acetylcholine receptor activation, membrane depolarization, or protein kinase C activation. This protein has been shown to bind CRK-associated substrate, nephrocystin, GTPase regulator associated with FAK, and the SH2 domain of GRB2. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Four transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human PYK2

Gene ID

2185

Swiss Prot

Q14289

Synonyms

PKB; PTK; CAKB; FAK2; PYK2; CADTK; FADK2; RAFTK

Reactivity

Human, Rat

Application

WB, IHC, IP

Recommended dilution

WB: 1:1000

IHC: 1:50

IP: 1:20

Calculated MW

116 kDa

Observed MW

116 kDa

Host species

Rabbit

Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.