



Ikaros Rabbit pAb

db21559 Package : 20μL 50μL 100μL

Product Name: Ikaros Rabbit pAb

Cat.No.: db21559

Synonyms: IK1; LYF1; LyF-1; CVID13; IKAROS; PPP1R92; PRO0758; ZNFN1A1; Hs.54452

Application: WB, ICC/IF, FC

Reactivity : Human

Host species : Rabbit

Background

This gene encodes a transcription factor that belongs to the family of zinc-finger DNA-binding proteins associated with chromatin remodeling. The expression of this protein is restricted to the fetal and adult hemo-lymphopoietic system, and it functions as a regulator of lymphocyte differentiation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. Most isoforms share a common C-terminal domain, which contains two zinc finger motifs that are required for hetero- or homo-dimerization, and for interactions with other proteins. The isoforms, however, differ in the number of N-terminal zinc finger motifs that bind DNA and in nuclear localization signal presence, resulting in members with and without DNA-binding properties. Only a few isoforms contain the requisite three or more N-terminal zinc motifs that confer high affinity binding to a specific core DNA sequence element in the promoters of target genes. The non-DNA-binding isoforms are largely found in the cytoplasm, and are thought to function as dominant-negative factors. Overexpression of some dominant-negative isoforms have been associated with B-cell malignancies, such as acute lymphoblastic leukemia (ALL). [provided

by RefSeq, May 2014]

Immunogen A synthetic peptide of human lkaros

Gene ID 10320

Swiss Prot Q13422

Synonyms IK1; LYF1; LyF-1; CVID13; IKAROS; PPP1R92; PR00758; ZNFN1A1; Hs.54452

Reactivity Human

Application WB, ICC/IF, FC

Recommended dilution WB: 1:1000

ICC/IF: 1:50

FC: 1:20

Calculated MW 58 kDa



For Research Use Only **Product Datasheet**

Observed MW 50-70 kDa

Host species Rabbit

Clonality Polyclonal

Isotype IgG

Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.