



Phospho-PKC zeta/lambda (Thr410/Thr412) Rabbit pAb

db21023 Package : 20μL 50μL 100μL

Product Name: Phospho-PKC zeta/lambda (Thr410/Thr412) Rabbit pAb

Cat.No.: db21023

Synonyms: PKCC; PKCG; SCA14; PKCI(3); PKCgamma; PKC-gamma

Application: WB, IHC

Reactivity: Human, Mouse, Rat

Host species: Rabbit

Background Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be

activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play distinct roles in cells.

expressed solely in the brain and spinal cord and its localization is restricted to neurons. It has been demonstrated that several neuronal functions, including long term potentiation (LTP) and long

The protein encoded by this gene is one of the PKC family members. This protein kinase is

term depression (LTD), specifically require this kinase. Knockout studies in mice also suggest that this kinase may be involved in neuropathic pain development. Defects in this protein have been associated with neurodegenerative disorder spinocerebellar ataxia-14 (SCA14). Two transcript

variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015]

Immunogen A synthetic phosphopeptide corresponding to residues surrounding Thr410 of human PKC zeta

Gene ID 5582, 5584, 5590

Swiss Prot P05129, P41743, Q05513

Synonyms PKCC; PKCG; SCA14; PKCl(3); PKCgamma; PKC-gamma

Reactivity Human, Mouse, Rat

Application WB, IHC

Recommended dilution WB: 1:1000

IHC: 1:20

Calculated MW 68 kDa

Observed MW 76 kDa

Host species Rabbit



For Research Use Only **Product Datasheet**

Clonality Polyclonal

Isotype IgG

Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.