

## Phospho-Histone H3 (Ser10/Thr11) Rabbit pAb

db21276

Package : 20µL 50µL 100µL

**Product Name** : Phospho-Histone H3 (Ser10/Thr11) Rabbit pAb**Cat.No.:** db21276**Synonyms** : H3/A; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FA; H3C10; H3C11; H3C12; HIST1H3A**Application** : WB, IHC, ICC/IF, IP**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015]

**Immunogen**

A synthetic phosphopeptide corresponding to residues surrounding Ser10/Thr11 of human Histone H3

**Gene ID**

8350

**Swiss Prot**

P68431

**Synonyms**

H3/A; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FA; H3C10; H3C11; H3C12; HIST1H3A

**Reactivity**

Human, Mouse, Rat

**Application**

WB, IHC, ICC/IF, IP

**Recommended dilution**

WB: 1:1000  
IHC: 1:20  
ICC/IF: 1:50-1:200  
IP: 1:20

**Calculated MW**

15 kDa

**Observed MW**

17 kDa

**Host species**

Rabbit

**Clonality**

Polyclonal

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.