

Phospho-FANCD2 (Ser1404) Rabbit pAb

db21963

Package : 20µL 50µL 100µL

Product Name : Phospho-FANCD2 (Ser1404) Rabbit pAb**Cat.No.:** db21963**Synonyms** : FA4; FAD; FACD; FAD2; FA-D2; FANCD**Application** : WB, ICC/IF**Reactivity** : Human, Mouse**Host species** : Rabbit**Background**

The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCI (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group D2. This protein is monoubiquitinated in response to DNA damage, resulting in its localization to nuclear foci with other proteins (BRCA1 AND BRCA2) involved in homology-directed DNA repair. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016]

Immunogen

A synthetic phosphopeptide corresponding to residues surrounding Ser1404 of human FANCD2

Gene ID

2177

Swiss Prot

Q9BXW9

Synonyms

FA4; FAD; FACD; FAD2; FA-D2; FANCD

Reactivity

Human, Mouse

Application

WB, ICC/IF

Recommended dilution

WB: 1:1000

ICC/IF: 1:50

Calculated MW

55 kDa

Observed MW

55 kDa

Host species

Rabbit

Clonality

Polyclonal

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.