Phospho-FANCD2 (Ser1404) Rabbit pAb

db21963 **Package : 20μL 50μL 100μL**

Product Name : Phospho-FANCD2 (Ser1404) Rabbit pAb
Cat.No.: db21963
Synonyms : FA4; FAD; FACD; FAD2; FA-D2; FANCD
Application : WB, ICC/IF
Reactivity : Human, Mouse
Host species : Rabbit

Background	The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCJ (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group D2. This protein is monoubiquinated in response to DNA damage, resulting in its localization to nuclear foci with other proteins (BRCA1 AND BRCA2) involved in homology-directed DNA repair. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016]
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Ser1404 of human FANCD2
Gene ID	2177
Swiss Prot	Q9BXW9
Synonyms	FA4; FAD; FACD; FAD2; FA-D2; FANCD
Reactivity	Human, Mouse
Application	WB, ICC/IF
Recommended dilution	WB: 1:1000 ICC/IF: 1:50
Calculated MW	55 kDa
Observed MW	55 kDa
Host species	Rabbit
Clonality	Polyclonal

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.