

## Bcl-XL Rabbit pAb

db225

Package : 20μL 50μL 100μL

**Product Name** : Bcl-XL Rabbit pAb**Cat.No.:** db225**Synonyms** : BCLX; BCL2L; Bcl-X; PPP1R52; BCL-XL/S**Application** : WB, IHC, ICC/IF, FC, IP**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Alternative splicing results in multiple transcript variants encoding two different isoforms. The longer isoform acts as an apoptotic inhibitor and the shorter isoform acts as an apoptotic activator. [provided by RefSeq, Dec 2015]

**Immunogen**

A synthetic peptide of human Bcl-XL

**Gene ID**

598

**Swiss Prot**

Q07817

**Synonyms**

BCLX; BCL2L; Bcl-X; PPP1R52; BCL-XL/S

**Reactivity**

Human, Mouse, Rat

**Application**

WB, IHC, ICC/IF, FC, IP

**Recommended dilution**WB: 1:1000  
IHC: 1:200  
ICC/IF: 1:100  
FC: 1:20  
IP: 1:30**Calculated MW**

26 kDa

**Observed MW**

30 kDa

**Host species**

Rabbit

<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.