

IL-4 Rabbit pAb**db398****Package : 20µL 50µL 100µL****Product Name :** IL-4 Rabbit pAb**Cat.No.:** db398**Synonyms :** BSF1; IL-4; BCGF1; BSF-1; BCGF-1**Application :** WB, ICC/IF, FC**Reactivity :** Human**Host species :** Rabbit**Background**

The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human IL4

Gene ID

3565

Swiss Prot

P05112

Synonyms

BSF1; IL-4; BCGF1; BSF-1; BCGF-1

Reactivity

Human

Application

WB, ICC/IF, FC

Recommended dilution

WB: 1:1000

ICC/IF: 1:50

FC: 1:20

Calculated MW

17 kDa

Observed MW

17 kDa

Host species

Rabbit

Clonality

Polyclonal

Isotype

IgG

| | |
|--------------------------|---|
| Purity | Affinity Purification |
| Conjugation | Un-conjugated |
| Storage Stability | Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. |

□