



NFkB p100 Rabbit pAb

db436 Package: 20μL 50μL 100μL

Product Name: NFkB p100 Rabbit pAb

Cat.No.: db436

Synonyms: p52; p100; H2TF1; LYT10; CVID10; LYT-10; NF-kB2; p49/p100

Application : WB, ICC/IF, FC, IP **Reactivity :** Human, Mouse, Rat

Host species: Rabbit

Background This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB).

The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on

which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants. [provided by RefSeq,

Dec 2013]

Immunogen Recombinant protein of human NFkB p100

Gene ID 4791

Swiss Prot Q00653

Synonyms p52; p100; H2TF1; LYT10; CVID10; LYT-10; NF-kB2; p49/p100

Reactivity Human, Mouse, Rat

Application WB, ICC/IF, FC, IP

Recommended dilution WB: 1:1000

ICC/IF: 1:50 FC: 1:20

IP: 1:20

Calculated MW 97 kDa

Observed MW 120 kDa

Host species Rabbit

Clonality Polyclonal



For Research Use Only **Product Datasheet**

Isotype IgG

Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.