

**SHP2 Rabbit pAb****db445****Package : 20µL 50µL 100µL****Product Name :** SHP2 Rabbit pAb**Cat.No.:** db445**Synonyms :** CFC; NS1; JMML; SHP2; BPTP3; PTP2C; METCDS; PTP-1D; SH-PTP2; SH-PTP3**Application :** WB, IHC, ICC/IF, FC, IP**Reactivity :** Human**Host species :** Rabbit**Background**

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains two tandem Src homology-2 domains, which function as phospho-tyrosine binding domains and mediate the interaction of this PTP with its substrates. This PTP is widely expressed in most tissues and plays a regulatory role in various cell signaling events that are important for a diversity of cell functions, such as mitogenic activation, metabolic control, transcription regulation, and cell migration. Mutations in this gene are a cause of Noonan syndrome as well as acute myeloid leukemia. [provided by RefSeq, Aug 2016]

**Immunogen**

A synthetic peptide of human SHP2

**Gene ID**

5781

**Swiss Prot**

Q06124

**Synonyms**

CFC; NS1; JMML; SHP2; BPTP3; PTP2C; METCDS; PTP-1D; SH-PTP2; SH-PTP3

**Reactivity**

Human

**Application**

WB, IHC, ICC/IF, FC, IP

**Recommended dilution**

WB: 1:2000

IHC: 1:20

ICC/IF: 1:20

FC: 1:20

IP: 1:20

**Calculated MW**

68 kDa

**Observed MW**

68 kDa

**Host species**

Rabbit

---

<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

□