

HIF-1-alpha Rabbit pAb

db489

Package : 20µL 50µL 100µL

Product Name : HIF-1-alpha Rabbit pAb**Cat.No.:** db489**Synonyms** : HIF1; MOP1; PASD8; HIF-1A; bHLHe78; HIF-1alpha; HIF1-ALPHA; HIF-1-alpha**Application** : WB, IHC, ICC/IF, FC, IP**Reactivity** : Human**Host species** : Rabbit**Background**

This gene encodes the alpha subunit of transcription factor hypoxia-inducible factor-1 (HIF-1), which is a heterodimer composed of an alpha and a beta subunit. HIF-1 functions as a master regulator of cellular and systemic homeostatic response to hypoxia by activating transcription of many genes, including those involved in energy metabolism, angiogenesis, apoptosis, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia. HIF-1 thus plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jul 2011]

Immunogen

A synthetic peptide of human HIF-1-alpha

Gene ID

3091

Swiss Prot

Q16665

Synonyms

HIF1; MOP1; PASD8; HIF-1A; bHLHe78; HIF-1alpha; HIF1-ALPHA; HIF-1-alpha

Reactivity

Human

Application

WB, IHC, ICC/IF, FC, IP

Recommended dilutionWB: 1:1000
IHC: 1:50
ICC/IF: 1:250
FC: 1:50
IP: 1:50**Calculated MW**

93 kDa

Observed MW

120 kDa

Host species

Rabbit

Clonality

Polyclonal

| | |
|--------------------------|---|
| Isotype | IgG |
| Purity | Affinity Purification |
| Conjugation | Un-conjugated |
| Storage Stability | Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. |

□