

CD13 Rabbit pAb

db694

Package : 20µL 50µL 100µL

Product Name : CD13 Rabbit pAb Cat.No.: db694 Synonyms : APN; CD13; LAP1; P150; PEPN; GP150 Application : WB, IHC, ICC/IF, IP Reactivity : Human, Rat Host species : Rabbit

| Background | Aminopeptidase N is located in the small-intestinal and renal microvillar membrane, and also in |
|----------------------|--|
| | other plasma membranes. In the small intestine aminopeptidase N plays a role in the final digestion |
| | of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Its function |
| | in proximal tubular epithelial cells and other cell types is less clear. The large extracellular |
| | carboxyterminal domain contains a pentapeptide consensus sequence characteristic of members |
| | of the zinc-binding metalloproteinase superfamily. Sequence comparisons with known enzymes of |
| | this class showed that CD13 and aminopeptidase N are identical. The latter enzyme was thought to |
| | be involved in the metabolism of regulatory peptides by diverse cell types, including small intestinal |
| | and renal tubular epithelial cells, macrophages, granulocytes, and synaptic membranes from the |
| | CNS. Human aminopeptidase N is a receptor for one strain of human coronavirus that is an |
| | important cause of upper respiratory tract infections. Defects in this gene appear to be a cause of |
| | various types of leukemia or lymphoma. [provided by RefSeq, Jul 2008] |
| Immunogen | A synthetic peptide of human CD13 |
| Gene ID | 290 |
| Swiss Prot | P15144 |
| Synonyms | APN; CD13; LAP1; P150; PEPN; GP150 |
| Reactivity | Human, Rat |
| Application | WB, IHC, ICC/IF, IP |
| Recommended dilution | WB: 1:1000-1:5000 |
| | IHC: 1:20-1:100 |
| | ICC/IF: 1:20-1:50 |
| | IP: 1:20-1:50 |
| Calculated MW | 110 kDa |
| Observed MW | 160 kDa |

dvagbvo 戴格生物

| Host species | Rabbit |
|-------------------|---|
| Clonality | Polyclonal |
| lsotype | lgG |
| Purity | Affinity Purification |
| Conjugation | Un-conjugated |
| Storage Stability | Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium |
| | azide and 0.05% BSA. Stable for 12 months from date of receipt. |