

## HIF-1-alpha Rabbit pAb

db794

Package : 20µL 50µL 100µL

Product Name : HIF-1-alpha Rabbit pAb Cat.No.: db794 Synonyms : HIF1; MOP1; PASD8; HIF-1A; bHLHe78; HIF-1alpha; HIF1-ALPHA; HIF-1-alpha Application : WB, ICC/IF Reactivity : Human, Mouse, Rat Host species : Rabbit

Background	This gene encodes the alpha subunit of transcription factor hypoxia-inducible factor-1 (HIF-1), which is a heterodimer composed of an alpha and a beta subunit. HIF-1 functions as a master regulator of cellular and systemic homeostatic response to hypoxia by activating transcription of many genes, including those involved in energy metabolism, angiogenesis, apoptosis, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia. HIF-1 thus plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jul 2011]
Immunogen	Recombinant protein of human HIF-1-alpha
Gene ID	3091
Swiss Prot	Q16665
Synonyms	HIF1; MOP1; PASD8; HIF-1A; bHLHe78; HIF-1alpha; HIF1-ALPHA; HIF-1-alpha
Reactivity	Human, Mouse, Rat
Application	WB, ICC/IF
Application	
Recommended dilution	WB: 1:1000 ICC/IF: 1:50
Recommended dilution	ICC/IF: 1:50
Recommended dilution	ICC/IF: 1:50 92 kDa
Recommended dilution Calculated MW Observed MW	ICC/IF: 1:50 92 kDa 120 kDa
Recommended dilution Calculated MW Observed MW Host species	ICC/IF: 1:50 92 kDa 120 kDa Rabbit



Conjugation

Un-conjugated

**Storage Stability** 

Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.