

## PSMA Rabbit pAb

db1634

Package : 20µL 50µL 100µL

**Product Name** : PSMA Rabbit pAb**Cat.No.:** db1634**Synonyms** : PSM; FGCP; FOLH; GCP2; PSMA; mGCP; GCPII; NAALAD1; NAALAdase**Application** : WB, IHC, FC**Reactivity** : Human**Host species** : Rabbit**Background**

This gene encodes a type II transmembrane glycoprotein belonging to the M28 peptidase family. The protein acts as a glutamate carboxypeptidase on different alternative substrates, including the nutrient folate and the neuropeptide N-acetyl-L-aspartyl-L-glutamate and is expressed in a number of tissues such as prostate, central and peripheral nervous system and kidney. A mutation in this gene may be associated with impaired intestinal absorption of dietary folates, resulting in low blood folate levels and consequent hyperhomocysteinemia. Expression of this protein in the brain may be involved in a number of pathological conditions associated with glutamate excitotoxicity. In the prostate the protein is up-regulated in cancerous cells and is used as an effective diagnostic and prognostic indicator of prostate cancer. This gene likely arose from a duplication event of a nearby chromosomal region. Alternative splicing gives rise to multiple transcript variants encoding several different isoforms. [provided by RefSeq, Jul 2010]

**Immunogen**

A synthetic peptide of human PSMA

**Gene ID**

2346

**Swiss Prot**

Q04609

**Synonyms**

PSM; FGCP; FOLH; GCP2; PSMA; mGCP; GCPII; NAALAD1; NAALAdase

**Reactivity**

Human

**Application**

WB, IHC, FC

**Recommended dilution**WB: 1:5000  
IHC: 1:50  
FC: 1:20**Calculated MW**

84 kDa

**Observed MW**

100 kDa

**Host species**

Rabbit

Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.