

**c-Myc Rabbit pAb**

db1667

Package : 20µL 50µL 100µL

**Product Name** : c-Myc Rabbit pAb**Cat.No.:** db1667**Synonyms** : MRTL; MYCC; c-Myc; bHLHe39**Application** : WB, IHC, ICC/IF, FC, IP**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

This gene is a proto-oncogene and encodes a nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. The encoded protein forms a heterodimer with the related transcription factor MAX. This complex binds to the E box DNA consensus sequence and regulates the transcription of specific target genes. Amplification of this gene is frequently observed in numerous human cancers. Translocations involving this gene are associated with Burkitt lymphoma and multiple myeloma in human patients. There is evidence to show that translation initiates both from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site, resulting in the production of two isoforms with distinct N-termini. [provided by RefSeq, Aug 2017]

**Immunogen**

A synthetic peptide of human c-Myc

**Gene ID**

4609

**Swiss Prot**

P01106

**Synonyms**

MRTL; MYCC; c-Myc; bHLHe39

**Reactivity**

Human, Mouse, Rat

**Application**

WB, IHC, ICC/IF, FC, IP

**Recommended dilution**WB: 1:2000  
IHC: 1:200  
ICC/IF: 1:100  
FC: 1:100  
IP: 1:50**Calculated MW**

49 kDa

**Observed MW**

57-65 kDa

**Host species**

Rabbit

Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.