



## MCM7 Rabbit pAb

db1729 Package: 20μL 50μL 100μL

Product Name: MCM7 Rabbit pAb

Cat.No.: db1729

Synonyms: MCM2; CDC47; P85MCM; P1CDC47; PNAS146; PPP1R104; P1.1-MCM3

Application: WB, ICC/IF, FC

Reactivity : Human

Host species : Rabbit

**Background** The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance

proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex (pre\_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1-dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumorsuppressor protein RB1/RB. Alternatively spliced transcript variants

encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

**Immunogen** A synthetic peptide of human MCM7

**Gene ID** 4176

Swiss Prot P33993

**Synonyms** MCM2; CDC47; P85MCM; P1CDC47; PNAS146; PPP1R104; P1.1-MCM3

Reactivity Human

**Application** WB, ICC/IF, FC

Recommended dilution WB: 1:1000

ICC/IF: 1:20-1:50

FC: 1:20

Calculated MW 81 kDa

Observed MW 81 kDa

Host species Rabbit

**Clonality** Polyclonal

**Isotype** IgG



## For Research Use Only **Product Datasheet**

**Purity** Affinity Purification

**Conjugation** Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.