

Insulin Rabbit pAb

db1813

Package : 20μL 50μL 100μL

Product Name : Insulin Rabbit pAb**Cat.No.:** db1813**Synonyms** : IDDM; ILPR; IRDN; IDDM1; IDDM2; MODY10**Application** : WB, IHC**Reactivity** : Human**Host species** : Rabbit**Background**

After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into three peptides: the B chain and A chain peptides, which are covalently linked via two disulfide bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR) stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2010]

Immunogen

A synthetic peptide of human Insulin

Gene ID

3630

Swiss Prot

P01308

Synonyms

IDDM; ILPR; IRDN; IDDM1; IDDM2; MODY10

Reactivity

Human

Application

WB, IHC

Recommended dilutionWB: 1:1000
IHC: 1:50**Calculated MW**

12 kDa

Observed MW

12 kDa

Host species

Rabbit

Clonality

Polyclonal

Isotype

IgG

Purity

Affinity Purification

Conjugation

Un-conjugated

Storage Stability

Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.