Insulin Rabbit pAb

db1813 **Package : 20μL 50μL 100μL**

Product Name : Insulin Rabbit pAb
Cat.No.: db1813
Synonyms : IDDM; ILPR; IRDN; IDDM1; IDDM2; MODY10
Application : WB, IHC
Reactivity : Human
Host species : Rabbit

Background	After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into three peptides: the B chain and A chain peptides, which are covalently linked via two disulfide bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR) stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2010]
Immunogen	A synthetic peptide of human Insulin
Gene ID	3630
Swiss Prot	P01308
Synonyms	IDDM; ILPR; IRDN; IDDM1; IDDM2; MODY10
Reactivity	Human
Application	WB, IHC
Recommended dilution	WB: 1:1000 IHC: 1:50
Calculated MW	12 kDa
Observed MW	12 kDa
Host species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated

Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.