

Notch1 Rabbit pAb

db1840

Package : 20μL 50μL 100μL

Product Name : Notch1 Rabbit pAb**Cat.No.:** db1840**Synonyms** : hN1; AOS5; TAN1; AOVD1**Application** : WB, IHC, ICC/IF, FC**Reactivity** : Human, Mouse**Host species** : Rabbit**Background**

This gene encodes a member of the NOTCH family of proteins. Members of this Type I transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple different domain types. Notch signaling is an evolutionarily conserved intercellular signaling pathway that regulates interactions between physically adjacent cells through binding of Notch family receptors to their cognate ligands. The encoded preproprotein is proteolytically processed in the trans-Golgi network to generate two polypeptide chains that heterodimerize to form the mature cell-surface receptor. This receptor plays a role in the development of numerous cell and tissue types. Mutations in this gene are associated with aortic valve disease, Adams-Oliver syndrome, T-cell acute lymphoblastic leukemia, chronic lymphocytic leukemia, and head and neck squamous cell carcinoma. [provided by RefSeq, Jan 2016]

Immunogen

A synthetic peptide of human Notch1

Gene ID

4851

Swiss Prot

P46531

Synonyms

hN1; AOS5; TAN1; AOVD1

Reactivity

Human, Mouse

Application

WB, IHC, ICC/IF, FC

Recommended dilution

WB: 1:1000

IHC: 1:20

ICC/IF: 1:20

FC: 1:20

Calculated MW

273 kDa

Observed MW

120 kDa

Host species

Rabbit

Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.