



## Chk2 Rabbit pAb

db2417 Package: 20μL 50μL 100μL

Product Name: Chk2 Rabbit pAb

Cat.No.: db2417

Synonyms: CDS1; CHK2; LFS2; RAD53; hCds1; HuCds1; PP1425

Application: WB, IHC, ICC/IF, FC, IP

Reactivity : Human

Host species : Rabbit

## **Background**

In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage.

Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2012]

**Immunogen** Recombinant protein of human Chk2

**Gene ID** 11200

Swiss Prot 096017

**Synonyms** CDS1; CHK2; LFS2; RAD53; hCds1; HuCds1; PP1425

**Reactivity** Human

**Application** WB, IHC, ICC/IF, FC, IP

Recommended dilution WB: 1:5000-1:10000

IHC: 1:20-1:100 ICC/IF: 1:50 FC: 1:20

IP: 1:20-1:50

Calculated MW 61 kDa



## For Research Use Only **Product Datasheet**

**Observed MW** 61 kDa

Host species Rabbit

**Clonality** Polyclonal

**Isotype** IgG

**Purity** Affinity Purification

**Conjugation** Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.