

Cyclin E1 Rabbit pAb

db2420

Package : 20µL 50µL 100µL

Product Name : Cyclin E1 Rabbit pAb**Cat.No.:** db2420**Synonyms** : CCNE; pCCNE1**Application** : WB, ICC/IF**Reactivity** : Human**Host species** : Rabbit**Background**

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB. [provided by RefSeq, Apr 2016]

Immunogen

A synthetic peptide of human Cyclin E1

Gene ID

898

Swiss Prot

P24864

Synonyms

CCNE; pCCNE1

Reactivity

Human

Application

WB, ICC/IF

Recommended dilution

WB: 1:1000-1:5000

ICC/IF: 1:50

Calculated MW

47 kDa

Observed MW

47 kDa

Host species

Rabbit

Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.