

Cyclin H Rabbit pAb

db2597

Package : 20μL 50μL 100μL

Product Name : Cyclin H Rabbit pAb**Cat.No.:** db2597**Synonyms** : CAK; p34; p37; CycH**Application** : WB**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK7 kinase and ring finger protein MAT1. The kinase complex is able to phosphorylate CDK2 and CDC2 kinases, thus functions as a CDK-activating kinase (CAK). This cyclin and its kinase partner are components of TFIIH, as well as RNA polymerase II protein complexes. They participate in two different transcriptional regulation processes, suggesting an important link between basal transcription control and the cell cycle machinery. A pseudogene of this gene is found on chromosome 4. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Nov 2010]

Immunogen

A synthetic peptide of human Cyclin H

Gene ID

902

Swiss Prot

P51946

Synonyms

CAK; p34; p37; CycH

Reactivity

Human, Mouse, Rat

Application

WB

Recommended dilution

WB: 1:1000-1:5000

Calculated MW

38 kDa

Observed MW

36 kDa

Host species

Rabbit

Clonality

Polyclonal

Isotype

IgG

Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.