

## PTEN Rabbit pAb

db3274

Package : 20µL 50µL 100µL

Product Name : PTEN Rabbit pAb Cat.No.: db3274 Synonyms : BZS; DEC; CWS1; GLM2; MHAM; TEP1; MMAC1; PTEN1; 10q23del; PTENbeta Application : WB, IHC, ICC/IF, FC Reactivity : Human, Mouse, Rat Host species : Rabbit

Background	This gene was identified as a tumor suppressor that is mutated in a large number of cancers at
	high frequency. The protein encoded by this gene is a phosphatidylinositol-3,4,5-trisphosphate 3-
	phosphatase. It contains a tensin like domain as well as a catalytic domain similar to that of the
	dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases,
	this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates
	intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor
	suppressor by negatively regulating AKT/PKB signaling pathway. The use of a non-canonical
	(CUG) upstream initiation site produces a longer isoform that initiates translation with a leucine,
	and is thought to be preferentially associated with the mitochondrial inner membrane. This longer
	isoform may help regulate energy metabolism in the mitochondria. A pseudogene of this gene is
	found on chromosome 9. Alternative splicing and the use of multiple translation start codons results
	in multiple transcript variants encoding different isoforms. [provided by RefSeq, Feb 2015]
Immunogen	Recombinant protein of human PTEN
Gene ID	5728
Swiss Prot	P60484
Synonyms	BZS; DEC; CWS1; GLM2; MHAM; TEP1; MMAC1; PTEN1; 10q23del; PTENbeta
Reactivity	Human, Mouse, Rat
Application	WB, IHC, ICC/IF, FC
Recommended dilution	WB: 1:1000
	IHC: 1:20
	ICC/IF: 1:100
	FC: 1:20
Calculated MW	47 kDa
Observed MW	54 kDa

## dvagbvo 戴格生物

nal
Purification
ugated
t -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium nd 0.05% BSA. Stable for 12 months from date of receipt.
j