



Activin Receptor Type IA Rabbit pAb

db3891 Package: 20μL 50μL 100μL

Product Name: Activin Receptor Type IA Rabbit pAb

Cat.No.: db3891

Synonyms: FOP; ALK2; SKR1; TSRI; ACTRI; ACVR1A; ACVRLK2

Application: WB, IP

Reactivity: Human, Mouse, Rat

Host species: Rabbit

Background Activins are dimeric growth and differentiation factors which belong to the transforming growth

factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. This gene encodes activin A type I receptor

which signals a particular transcriptional response in concert with activin type II receptors.

Mutations in this gene are associated with fibrodysplasia ossificans progressive. [provided by

RefSeq, Jul 2008]

Immunogen A synthetic peptide of human Activin Receptor Type IA

Gene ID 90

Swiss Prot Q04771

Synonyms FOP; ALK2; SKR1; TSRI; ACTRI; ACVR1A; ACVRLK2

Reactivity Human, Mouse, Rat

Application WB, IP

Recommended dilution WB: 1:1000-1:5000

IP: 1:20

Calculated MW 57 kDa

Observed MW 57 kDa

Host species Rabbit



For Research Use Only **Product Datasheet**

Clonality Polyclonal

Isotype IgG

Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.