

p21 Rabbit pAb

db3999

Package : 20µL 50µL 100µL

Product Name : p21 Rabbit pAb**Cat.No.:** db3999**Synonyms** : P21; CDKI; CIP1; SDI1; Waf1; mda6; CAP20; Cdkn1; p21WAF; p21Cip1**Application** : WB, IHC, ICC/IF, FC, IP**Reactivity** : Mouse**Host species** : Rabbit**Background**

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at the G1 pahse. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]

Immunogen

Recombinant protein of mouse p21

Gene ID

12575

Swiss Prot

P39689

Synonyms

P21; CDKI; CIP1; SDI1; Waf1; mda6; CAP20; Cdkn1; p21WAF; p21Cip1

Reactivity

Mouse

Application

WB, IHC, ICC/IF, FC, IP

Recommended dilutionWB: 1:1000
IHC: 1:200
ICC/IF: 1:100
FC: 1:20
IP: 1:20**Calculated MW**

18 kDa

Observed MW

18 kDa

Host species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.