

## Raf1 Rabbit pAb

db4640

Package : 20μL 50μL 100μL

**Product Name** : Raf1 Rabbit pAb**Cat.No.:** db4640**Synonyms** : NS5; CRAF; Raf-1; c-Raf; CMD1NN**Application** : WB, ICC/IF, FC**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

This gene is the cellular homolog of viral raf gene (v-raf). The encoded protein is a MAP kinase kinase kinase (MAP3K), which functions downstream of the Ras family of membrane associated GTPases to which it binds directly. Once activated, the cellular RAF1 protein can phosphorylate to activate the dual specificity protein kinases MEK1 and MEK2, which in turn phosphorylate to activate the serine/threonine specific protein kinases, ERK1 and ERK2. Activated ERKs are pleiotropic effectors of cell physiology and play an important role in the control of gene expression involved in the cell division cycle, apoptosis, cell differentiation and cell migration. Mutations in this gene are associated with Noonan syndrome 5 and LEOPARD syndrome 2. [provided by RefSeq, Jul 2008]

**Immunogen**

A synthetic peptide of human Raf1

**Gene ID**

5894

**Swiss Prot**

P04049

**Synonyms**

NS5; CRAF; Raf-1; c-Raf; CMD1NN

**Reactivity**

Human,Mouse,Rat

**Application**

WB, ICC/IF, FC

**Recommended dilution**WB: 1:1000  
ICC/IF: 1:100  
FC: 1:50**Calculated MW**

73 kDa

**Observed MW**

73 kDa

**Host species**

Rabbit

**Clonality**

Polyclonal

**Isotype**

IgG

<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.