

IL-12A Rabbit pAb

db5107

Package : 20µL 50µL 100µL

Product Name : IL-12A Rabbit pAb Cat.No.: db5107 Synonyms : P35; CLMF; NFSK; NKSF1; IL-12A Application : WB Reactivity : Human, Mouse Host species : Rabbit

Background	This gene encodes a subunit of a cytokine that acts on T and natural killer cell array of biological activities. The cytokine is a disulfide-linked heterodimer co subunit encoded by this gene, and a 40-kD subunit that is a member of the cy family. This cytokine is required for the T-cell-independent induction of interfer is important for the differentiation of both Th1 and Th2 cells. The responses of cytokine are mediated by the activator of transcription protein STAT4. Nitric of (NOS2A/NOS2) is found to be required for the signaling process of this cytok [provided by RefSeq, Jul 2008]	mposed of the 35-kD tokine receptor on (IFN)-gamma, and ^r lymphocytes to this xide synthase 2A
Immunogen	A synthetic peptide of human IL12A	
Gene ID	3592	
Swiss Prot	P29459	
Synonyms	P35; CLMF; NFSK; NKSF1; IL-12A	
Reactivity	Human, Mouse	
Application	WB	
Recommended dilution	WB: 1:1000-1:5000	
Calculated MW	25 kDa	
Observed MW	35 kDa	
Host species	Rabbit	
Clonality	Polyclonal	
lsotype	lgG	
Purity	Affinity Purification	
Conjugation	Un-conjugated	
www.diagbio.com	service@diagbio.com	0571 - 87000691



Storage Stability

Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.