

Smad4 Rabbit pAb

db5616

Package : 20μL 50μL 100μL

Product Name : Smad4 Rabbit pAb**Cat.No.:** db5616**Synonyms** : JIP; DPC4; MADH4; MYHRS**Application** : WB, IHC**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

This gene encodes a member of the Smad family of signal transduction proteins. Smad proteins are phosphorylated and activated by transmembrane serine-threonine receptor kinases in response to transforming growth factor (TGF)-beta signaling. The product of this gene forms homomeric complexes and heteromeric complexes with other activated Smad proteins, which then accumulate in the nucleus and regulate the transcription of target genes. This protein binds to DNA and recognizes an 8-bp palindromic sequence (GTCTAGAC) called the Smad-binding element (SBE). The protein acts as a tumor suppressor and inhibits epithelial cell proliferation. It may also have an inhibitory effect on tumors by reducing angiogenesis and increasing blood vessel hyperpermeability. The encoded protein is a crucial component of the bone morphogenetic protein signaling pathway. The Smad proteins are subject to complex regulation by post-translational modifications. Mutations or deletions in this gene have been shown to result in pancreatic cancer, juvenile polyposis syndrome, and hereditary hemorrhagic telangiectasia syndrome. [provided by RefSeq, Aug 2017]

Immunogen

A synthetic peptide of human Smad4

Gene ID

4089

Swiss Prot

Q13485

Synonyms

JIP; DPC4; MADH4; MYHRS

Reactivity

Human, Mouse, Rat

Application

WB, IHC

Recommended dilution

WB: 1:2000

IHC: 1:100

Calculated MW

60 kDa

Observed MW

70 kDa

Host species

Rabbit

Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

□