

Tissue Factor Rabbit pAb

db5881

Package : 20µL 50µL 100µL

Product Name : Tissue Factor Rabbit pAb**Cat.No.:** db5881**Synonyms** : TF; TFA; CD142**Application** : WB, IHC**Reactivity** : Human**Host species** : Rabbit**Background**

This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.[provided by RefSeq, May 2010]

Immunogen

Recombinant protein of human Tissue Factor

Gene ID

2152

Swiss Prot

P13726

Synonyms

TF; TFA; CD142

Reactivity

Human

Application

WB, IHC

Recommended dilution

WB: 1:1000

IHC: 1:20

Calculated MW

33 kDa

Observed MW

45-53 kDa

Host species

Rabbit

Clonality

Polyclonal

Isotype

IgG

Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.