

## Lamin A/C Rabbit pAb

db5914

Package : 20μL 50μL 100μL

**Product Name** : Lamin A/C Rabbit pAb

**Cat.No.:** db5914

**Synonyms** : FPL; IDC; LFP; CDDC; EMD2; FPLD; HGPS; LDP1; LMN1; LMNC; MADA; PRO1; CDCD1; CMD1A; FPLD2; LMNL1; CMT2B1; LGMD1B

**Application** : WB, IHC, ICC/IF, FC, IP

**Reactivity** : Human, Mouse, Rat

**Host species** : Rabbit

### Background

The nuclear lamina consists of a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Lamin proteins are thought to be involved in nuclear stability, chromatin structure and gene expression. Vertebrate lamins consist of two types, A and B. Alternative splicing results in multiple transcript variants. Mutations in this gene lead to several diseases: Emery-Dreifuss muscular dystrophy, familial partial lipodystrophy, limb girdle muscular dystrophy, dilated cardiomyopathy, Charcot-Marie-Tooth disease, and Hutchinson-Gilford progeria syndrome. [provided by RefSeq, Apr 2012]

### Immunogen

A synthetic peptide of human Lamin A/C

### Gene ID

4000

### Swiss Prot

P02545

### Synonyms

FPL; IDC; LFP; CDDC; EMD2; FPLD; HGPS; LDP1; LMN1; LMNC; MADA; PRO1; CDCD1; CMD1A; FPLD2; LMNL1; CMT2B1; LGMD1B

### Reactivity

Human, Mouse, Rat

### Application

WB, IHC, ICC/IF, FC, IP

### Recommended dilution

WB: 1:1000  
IHC: 1:20  
ICC/IF: 1:20  
FC: 1:20  
IP: 1:20

### Calculated MW

74 kDa

### Observed MW

74,63 kDa

<b>Host species</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.