# MEKK2 Rabbit pAb

**db7008** **Package : 20μL 50μL 100μL**

**Product Name :** MEKK2 Rabbit pAb
**Cat.No.:** db7008
**Synonyms :** MEKK2; MEKK2B
**Application :** WB, IHC, ICC/IF, FC, IP
**Reactivity :** Human, Mouse, Rat
**Host species :** Rabbit

| | |
|---|---|
| **Background** | The protein encoded by this gene is a member of serine/threonine protein kinase family. This kinase preferentially activates other kinases involved in the MAP kinase signaling pathway. This kinase has been shown to directly phosphorylate and activate Ikappa B kinases, and thus plays a role in NF-kappa B signaling pathway. This kinase has also been found to bind and activate protein kinase C-related kinase 2, which suggests its involvement in a regulated signaling process. [provided by RefSeq, Jul 2008] |
| **Immunogen** | A synthetic peptide of human MEKK2 |
| **Gene ID** | 10746 |
| **Swiss Prot** | Q9Y2U5 |
| **Synonyms** | MEKK2; MEKK2B |
| **Reactivity** | Human, Mouse, Rat |
| **Application** | WB, IHC, ICC/IF, FC, IP |
| **Recommended dilution** | WB: 1:2000-1:10000<br>IHC: 1:20<br>ICC/IF: 1:50<br>FC: 1:20<br>IP: 1:20 |
| **Calculated MW** | 70 kDa |
| **Observed MW** | 70 kDa |
| **Host species** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | IgG |

| | |
|---|---|
| **Purity** | Affinity Purification |
| **Conjugation** | Un-conjugated |
| **Storage Stability** | Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. |