



## Histone H2A.Z Rabbit pAb

db7330 Package: 20μL 50μL 100μL

Product Name: Histone H2A.Z Rabbit pAb

Cat.No.: db7330

**Synonyms:** H2AZ; H2A.z; H2A/z; H2A.Z-1

**Application:** WB, IHC, ICC/IF **Reactivity:** Human, Mouse, Rat

Host species: Rabbit

Background Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent member of the histone H2A family that is distinct from other members of the family. Studies in mice have shown that this particular histone is required for embryonic

development and indicate that lack of functional histone H2A leads to embryonic lethality. [provided

by RefSeq, Jul 2008]

**Immunogen** A synthetic peptide of human Histone H2A.Z

**Gene ID** 3015

Swiss Prot P0C0S5

**Synonyms** H2AZ; H2A.z; H2A/z; H2A.Z-1

Reactivity Human, Mouse, Rat

Application WB, IHC, ICC/IF

Recommended dilution WB: 1:1000

IHC: 1:50 ICC/IF: 1:50

ICC/IF: 1:50

Calculated MW 14 kDa

**Observed MW** 14 kDa

Host species Rabbit

**Clonality** Polyclonal

**Isotype** IgG



## For Research Use Only **Product Datasheet**

**Purity** Affinity Purification

**Conjugation** Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.