



ERG Rabbit pAb

db7938 Package : 20μL 50μL 100μL

Product Name: ERG Rabbit pAb

Cat.No.: db7938

Synonyms: p55; erg-3
Application: WB, IHC
Reactivity: Human, Mouse
Host species: Rabbit

Background

This gene encodes a member of the erythroblast transformation-specific (ETS) family of transcriptions factors. All members of this family are key regulators of embryonic development, cell proliferation, differentiation, angiogenesis, inflammation, and apoptosis. The protein encoded by this gene is mainly expressed in the nucleus. It contains an ETS DNA-binding domain and a PNT (pointed) domain which is implicated in the self-association of chimeric oncoproteins. This protein is required for platelet adhesion to the subendothelium, inducing vascular cell remodeling. It also regulates hematopoesis, and the differentiation and maturation of megakaryocytic cells. This gene is involved in chromosomal translocations, resulting in different fusion gene products, such as TMPSSR2-ERG and NDRG1-ERG in prostate cancer, EWS-ERG in Ewing's sarcoma and FUS-ERG in acute myeloid leukemia. More than two dozens of transcript variants generated from combinatorial usage of three alternative promoters and multiple alternative splicing events have been reported, but the full-length nature of many of these variants has not been determined. [provided by RefSeq, Apr 2014]

Immunogen A synthetic peptide of human ERG

Gene ID 2078

Swiss Prot P11308

Synonyms p55; erg-3

Reactivity Human, Mouse

Application WB, IHC

Recommended dilution WB: 1:1000

IHC: 1:50

Calculated MW 55 kDa

Observed MW 55 kDa

Host species Rabbit



For Research Use Only **Product Datasheet**

Clonality Polyclonal

Isotype IgG

Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.